

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C23G 5/028, C11D 7/50	A2	(11) International Publication Number: WO 92/00402 (43) International Publication Date: 9 January 1992 (09.01.92)
(21) International Application Number: PCT/US91/04521 (22) International Filing Date: 24 June 1991 (24.06.91) (30) Priority data: 543,238 25 June 1990 (25.06.90) US (71) Applicant: ALLIED-SIGNAL INC. [US/US]; Law Department (C.A. McNally), P.O. Box 2245R, Morristown, NJ 07962-2245 (US). (72) Inventors: MAGID, Hillel ; 19 Rockford Place, Buffalo, NY 14221 (US). WILSON, David, Paul ; 118 Waxwing Court, East Amherst, NY 14051 (US). HOLLISTER, Richard, M. ; 221 Englewood Avenue, Buffalo, NY 14214 (US). LAVERY, Dennis, M. ; 10210 Boston State Road, Springville, NY 14141 (US).		(74) Agent: ROONEY, Gerard, P.; Allied-Signal Inc., Law Department (C.A. McNally), P.O. Box 2245, Morristown, NJ 07962-2245 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), CS, DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent), SU. Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: AZEOTROPE-LIKE COMPOSITIONS OF DICHLOROPENTAFLUOROPROPANE, 2-PROPANOL AND A HYDROCARBON CONTAINING SIX CARBON ATOMS (57) Abstract <p>Novel azeotrope-like compositions comprising dichloropentafluoropropane, 2-propanol and a hydrocarbon containing six carbon atoms which are useful in a variety of industrial cleaning applications including cold cleaning and defluxing of printed circuit boards.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LJ	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

-1-

AZEOTROPE-LIKE COMPOSITIONS OF
DICHLOROPENTAFLUOROPROPANE, 2-PROPANOL AND
A HYDROCARBON CONTAINING SIX CARBON ATOMS

5

Field of the Invention

This invention relates to azeotrope-like mixtures of dichloropentafluoropropane, 2-propanol and a
10 hydrocarbon containing six carbon atoms. These mixtures are useful in a variety of vapor degreasing, cold cleaning, and solvent cleaning applications including defluxing and dry cleaning.

15

CROSS-REFERENCE TO RELATED APPLICATIONS

Co-pending commonly assigned patent application serial no.: 418,008, filed October 6, 1989, discloses azeotrope-like compositions of 1,1-dichloro-2,2,3,3,3-
20 pentafluoropropane and alkanol having 1 to 3 carbon atoms.

Co-pending commonly assigned patent application serial no.: 417,983, filed October 6, 1989, discloses azeotrope-like compositions of 1,3-dichloro-1,1,2,2,3-
25 pentafluoropropane and alkanol having 1 to 3 carbon atoms.

Co-pending commonly assigned patent application serial no.: 526,748, filed May 22, 1990, discloses azeotrope-like compositions of dichloropentafluoropropane and alkanol having 1 to 4 carbon atoms.
30

Co-pending commonly assigned patent application Serial No.: 418,050, filed October 6, 1989, discloses azeotrope-like compositions of 1,1-dichloro-2,2,3,3,3-
35 pentafluoropropane and alkane having 6 carbon atoms.

-2-

Co-pending commonly assigned patent application
Serial No.: 417,951, filed October 6, 1989, discloses
azeotrope-like mixtures of 1,3-dichloro-1,1,2,2,3,3-
pentafluoropropane and cyclohexane.

Co-pending commonly assigned patent application
Serial No.: 454,789, filed December 21, 1989, discloses
azeotrope-like compositions of dichloropentafluoropropane
and cyclohexane.

Co-pending commonly assigned patent application
Serial No.: 526,874, filed May 22, 1990, discloses
azeotrope-like compositions of dichloropentafluoropropane
and a hydrocarbon containing six carbon atoms.

BACKGROUND OF THE INVENTION

Fluorocarbon based solvents have been used
extensively for the degreasing and otherwise cleaning of
solid surfaces, especially intricate parts and difficult
to remove soils.

In its simplest form, vapor degreasing or solvent
cleaning consists of exposing a room temperature object to
be cleaned to the vapors of a boiling solvent. Vapors
condensing on the object provide clean distilled solvent
to wash away grease or other contamination. Final
evaporation of solvent leaves the object of residue. This
is contrasted with liquid solvents which leave deposits on
the object after rinsing.

A vapor degreaser is used for difficult to remove
soils where elevated temperature is necessary to improve
the cleaning action of the solvent, or for large volume
assembly line operations where the cleaning of metal parts
and assemblies must be done efficiently. The conventional
operation of a vapor degreaser consists of immersing the

-3-

part to be cleaned in a sump of boiling solvent which removes the bulk of the soil, thereafter immersing the part in a sump containing freshly distilled solvent near room temperature, and finally exposing the part to solvent vapors over the boiling sump which condense on the cleaned part. In addition, the part can also be sprayed with distilled solvent before final rinsing.

Vapor degreasers suitable in the above-described operations are well known in the art. For example, Sherliker et al. in U.S. Patent 3,085,918 disclose such suitable vapor degreasers comprising a boiling sump, a clean sump, a water separator, and other ancillary equipment.

Cold cleaning is another application where a number of solvents are used. In most cold cleaning applications, the soiled part is either immersed in the fluid or wiped with cloths soaked in solvents and allowed to air dry.

Recently, non-toxic, non-flammable fluorocarbon solvents like trichlorotrifluoroethane, have been used extensively in degreasing applications and other solvent cleaning applications. Trichlorotrifluoroethane has been found to have satisfactory solvent power for greases, oils, waxes and the like. It has therefore found widespread use for cleaning electric motors, compressors, heavy metal parts, delicate precision metal parts, printed circuit boards, gyroscopes, guidance systems, aerospace and missile hardware, aluminum parts, etc.

The art has looked towards azeotropic compositions having fluorocarbon components because the fluorocarbon components contribute additionally desired characteristics, like polar functionality, increased solvency power, and stabilizers. Azeotropic compositions are desired because they do not fractionate upon boiling.

-4-

This behavior is desirable because in the previously described vapor degreasing equipment with which these solvents are employed, redistilled material is generated for final rinse-cleaning. Thus, the vapor degreasing system acts as a still. Therefore, unless the solvent composition is essentially constant boiling, fractionation will occur and undesirable solvent distribution may act to upset the cleaning and safety of processing. Preferential evaporation of the more volatile components of the solvent mixtures, which would be the case if they were not an azeotrope or azeotrope-like, would result in mixtures with changed compositions which may have less desirable properties, such as lower solvency towards soils, less inertness towards metal, plastic or elastomer components, and increased flammability and toxicity.

The art is continually seeking new fluorocarbon based azeotropic mixtures or azeotrope-like mixtures which offer alternatives for new and special applications for vapor degreasing and other cleaning applications. Currently, fluorocarbon-based azeotrope-like mixtures are of particular interest because they are considered to be stratospherically safe substitutes for presently used fully halogenated chlorofluorocarbons. The latter have been implicated in causing environmental problems associated with the depletion of the earth's protective ozone layer. Mathematical models have substantiated that hydrochlorofluorocarbons, like dichloropentafluoropropane, have a much lower ozone depletion potential and global warming potential than the fully halogenated species.

Accordingly, it is an object of this invention to provide novel environmentally acceptable azeotrope-like compositions based on dichloropentafluoropropane, ethanol and n-hexane which are useful in a variety of industrial cleaning applications.

-5-

It is another object of this invention to provide azeotrope-like compositions which are liquid at room temperature and will not fractionate under conditions of use.

Other objects and advantages of the invention will become apparent from the following description.

10 SUMMARY OF THE INVENTION

The invention relates to novel azeotrope-like compositions which are useful in a variety of industrial cleaning applications. Specifically, the invention relates to compositions of dichloropentafluoropropane, 2-propanol and a hydrocarbon containing six carbon atoms which are essentially constant boiling, environmentally acceptable and which remain liquid at room temperature.

20 DETAILED DESCRIPTION OF THE INVENTION

In accordance with the invention, novel azeotrope-like compositions have been discovered comprising from about 62 to about 99.8 weight percent dichloropentafluoropropane, from about 0.1 to about 10 weight percent 2-propanol and from about 0.1 to about 28 weight percent of a hydrocarbon containing six carbon atoms (HEREINAFTER referred to as "C₆ hydrocarbon") which boil at about 52.2°C ± about 3.9°C and preferably boil at about 52.4°C ± about 3.7°C at 760 mm Hg.

As used herein, the term "C₆ hydrocarbon" shall refer to aliphatic hydrocarbons having the empirical formula C₆H₁₄ and cycloaliphatic or substituted cycloaliphatic hydrocarbons having the empirical formula C₆H₁₂; and mixtures thereof.

-6-

Preferably, the term "C₆ hydrocarbon" refers to the following subset which includes: n-hexane, 2-methylpentane, 3-methylpentane, 2,2-dimethylbutane, 2,3-dimethylbutane, cyclohexane, methylcyclopentane, commercial isohexane (typically, the percentages of the isomers in commercial isohexane will fall into one of the two following formulations designated grade 1 and grade 2: grade 1: 35-75 weight percent 2-methylpentane, 10-40 weight percent 3-methylpentane, 7-30 weight percent 2,3-dimethylbutane, 7-30 weight percent 2,2-dimethylbutane, and 0.1-10.0 weight percent n-hexane, and up to about 5 weight percent other alkane isomers; the sum of the branched chain six carbon alkane isomers is about 90 to about 100 weight percent and the sum of the branched and straight chain six carbon alkane isomers is about 95 to about 100 weight percent; grade 2: 40-55 weight percent 2-methylpentane, 15-30 weight percent 3-methylpentane, 10-22 weight percent 2,3-dimethylbutane, 9-16 weight percent 2,2-dimethylbutane, and 0.1-5 weight percent n-hexane; the sum of the branched chain six carbon alkane isomers is about 95 to about 100 weight percent and the sum of the branched and straight chain six carbon alkane isomers is about 97 to about 100 weight percent) and mixtures thereof.

Dichloropentafluoropropane exists in nine isomeric forms: (1) 2,2-dichloro-1,1,1,3,3-pentafluoropropane (HCFC-225a); (2) 1,2-dichloro-1,2,3,3,3-pentafluoropropane (HCFC-225ba); (3) 1,2-dichloro-1,1,2,3,3-pentafluoropropane (HCFC-225bb); (4) 1,1-dichloro-2,2,3,3,3-pentafluoropropane (HCFC-225ca); (5) 1,3-dichloro-1,1,2,2,3-pentafluoropropane (HCFC-225cb); (6) 1,1-dichloro-1,2,2,3,3-pentafluoropropane (HCFC-225cc); (7) 1,2-dichloro-1,1,3,3,3-pentafluoropropane (HCFC-225d); (8) 1,3-dichloro-1,1,2,3,3-pentafluoropropane (HCFC-225ea); and (9) 1,1-dichloro-1,2,3,3,3-pentafluoropropane (HCFC-

-7-

225eb). For purposes of this invention, dichloropentafluoropropane will refer to any of the isomers or admixtures of the isomers in any proportion. The
5 1,1-dichloro-2,2,3,3,3-pentafluoropropane and
1,3-dichloro-1,1,2,2,3-pentafluoropropane isomers,
however, are the preferred isomers.

The dichloropentafluoropropane component of the
10 invention has good solvent properties. 2-propanol and the
hydrocarbon component are also good solvents. 2-propanol
dissolves polar organic materials and amine hydrochlorides
while the hydrocarbon enhances the solubility of oils.
Thus, when these components are combined in effective
15 amounts, an efficient azeotropic solvent results.

Preferably, the azeotrope like compositions consist
essentially of from about 67 to about 96.5 weight percent
dichloropentafluoropropane, from about 0.5 to about 7
20 weight percent 2-propanol and from about 3 to about 26
weight percent C₆ hydrocarbon.

In a more preferred embodiment of the invention,
the azeotrope-like compositions consist essentially of
25 from about 73 to about 95.5 weight percent dichloro-
pentafluoropropane, from about 0.5 to about 7 weight
percent 2-propanol and from about 4 to about 20 weight
percent C₆ hydrocarbon.

In another preferred embodiment of the invention,
30 the azeotrope-like compositions consist essentially of
from about 83 to about 95.5 weight percent dichloropenta-
fluoropropane, from about 0.5 to about 7 weight percent
2-propanol and from about 4 to about 10 weight percent
35 C₆ hydrocarbon.

In another preferred embodiment of the invention,
the azeotrope-like compositions consist essentially of

-8-

from about 67 to about 89.5 weight percent dichloro-
pentafluoropropane, from about 0.5 to about 7 weight
percent 2-propanol and from about 10 to about 26 weight
5 percent C_6 hydrocarbon.

When the C_6 hydrocarbon is n-hexane, the
azeotrope-like compositions of the invention consist
essentially of from about 73 to about 96.5 weight percent
10 dichloropentafluoropropane, from about 0.5 to about 7
weight percent 2-propanol and from about 3 to about 20
weight percent n-hexane and boil at about $52.6^{\circ}\text{C} \pm$ about
3.0 $^{\circ}\text{C}$ at 760 mm Hg.

When the C_6 hydrocarbon is 2-methylpentane, the
15 azeotrope-like compositions of the invention consist
essentially of from about 67 to about 93.5 weight percent
dichloropentafluoropropane, from about 0.5 to about 7
weight percent 2-propanol and from about 6 to about 26
weight percent 2-methylpentane and boil at about $51.4^{\circ}\text{C} \pm$
20 about 2.5 $^{\circ}\text{C}$ at 760 mm Hg.

When the C_6 hydrocarbon is 3-methylpentane, the
azeotrope-like compositions of the invention consist
25 essentially of from about 71 to about 96.5 weight percent
dichloropentafluoropropane, from about 0.5 to about 7
weight percent 2-propanol and from about 3 to about 22
weight percent 3-methylpentane and boil at about $51.9^{\circ}\text{C} \pm$
about 2.9 $^{\circ}\text{C}$ at 760 mm Hg.

30 When the C_6 hydrocarbon is methylcyclopentane,
the azeotrope-like compositions of the invention consist
essentially of from about 71 to about 96.5 weight percent
dichloropentafluoropropane, from about 0.5 to about 7
weight percent 2-propanol and from about 3 to about 22
35 weight percent methylcyclopentane and boil at about 52.8°C
 \pm about 2.8 $^{\circ}\text{C}$ at 760 mm Hg.

-9-

When the C₆ hydrocarbon is commercial isohexane grade 1, the azeotrope-like compositions of the invention consist essentially of from about 67 to about 93.5 weight percent of dichloropentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 6 to about 26 weight percent commercial isohexane grade 1 and boil at about 51.1°C ± about 2.7°C and preferably ± about 2.3°C at 760 mm Hg.

10

When the C₆ hydrocarbon is commercial isohexane grade 2, the azeotrope-like compositions of the invention consist essentially of from about 67 to about 93.5 weight percent dichloropentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 6 to about 26 weight percent commercial isohexane grade 2 and boil at about 51.1°C ± about 2.7°C and preferably ± about 2.3°C at 760 mm Hg.

15

When the C₆ hydrocarbon is cyclohexane, the azeotrope-like compositions of the invention consist essentially of from about 83 to about 99.4 weight percent dichloropentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 0.1 to about 10 weight percent cyclohexane and boil at about 53.3°C ± about 2.8°C at 760 mm Hg.

20

25

When the dichloropentafluoropropane component is 1,1,-dichloro-2,2,3,3,3-pentafluoropropane (225ca) and the C₆ hydrocarbon is n-hexane, the azeotrope-like compositions of the invention consist essentially of from about 87 to about 99.8 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 4 weight percent 2-propanol and from about 0.1 to about 9 weight percent n-hexane and boil at about 51.2°C ± about 0.7°C and preferably ± 0.5°C at 760 mm Hg.

30

35

-10-

In a preferred embodiment of the invention utilizing 225ca and n-hexane, the azeotrope-like compositions of the invention consist essentially of from about 88.5 to about 99.8 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 2.5 weight percent 2-propanol and from about 0.1 to about 9 weight percent n-hexane.

In another preferred embodiment of the invention utilizing 225ca and n-hexane, the azeotrope-like compositions consist essentially of from about 88 to about 99.4 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 4 weight percent 2-propanol and from about 0.5 to about 8 weight percent n-hexane.

In a more preferred embodiment of the invention utilizing 225ca and n-hexane, the azeotrope-like compositions of the invention consist essentially of from about 89.5 to about 99.4 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 2.5 weight percent 2-propanol and from about 0.5 to about 8 weight percent n-hexane.

In a still more preferred embodiment of the invention utilizing 225ca and n-hexane, the azeotrope-like compositions consist essentially of from about 91 to about 99.4 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 2 weight percent 2-propanol and from about 0.5 to about 7 weight percent n-hexane.

In a most preferred embodiment of the invention utilizing 225ca and n-hexane, the azeotrope-like compositions consist essentially of from about 92 to about 98.9 weight percent 1,1-dichloro-2,2,3,3,3-

-11-

pentafluoropropane, from about 0.1 to about 2 weight percent 2-propanol and from about 1 to about 6 weight percent n-hexane.

5

When the dichloropentafluoropropane component is 225ca and the C₆ hydrocarbon is 2-methylpentane, the azeotrope-like compositions of the invention consist essentially of from about 70 to about 92.5 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 7 to about 23 weight percent 2-methylpentane and boil at about 49.7°C ± about 0.9°C and preferably ± 0.5°C at 760 mm Hg.

10

15

When the dichloropentafluoropropane component is 225ca and the C₆ hydrocarbon is commercial isohexane grade 1, the azeotrope-like compositions of the invention consist essentially of from about 68 to about 92.5 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 7 to about 25 weight percent commercial isohexane grade 1 and boil at about 49.6°C ± about 1.3°C at 760 mm Hg.

20

25

When the dichloropentafluoropropane component is 225ca and the C₆ hydrocarbon is the commercial isohexane grade 2, the azeotrope-like compositions of the invention consist essentially of from about 68 to about 92.5 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 7 to about 25 weight percent commercial isohexane grade 2, and boil at about 49.6°C ± about 1.3°C at 760 mm Hg.

30

35

When the dichloropentafluoropropane component is 1,3-dichloro-1,1,2,2,3-pentafluoropropane (225cb) and the C₆ hydrocarbon is cyclohexane, the azeotrope-like compositions of the invention consist essentially of from

-12-

about 82 to about 99.4 weight percent 1,3-dichloro-
1,1,2,2,3-pentafluoropropane, from about 0.5 to about 9
weight percent 2-propanol and from about 0.1 to about 9
5 weight percent cyclohexane and boil at about $55.6^{\circ}\text{C} \pm$
about 0.5°C and preferably $\pm 0.2^{\circ}\text{C}$ at 760 mm Hg.

In a preferred embodiment of the invention
utilizing 225cb and cyclohexane, the azeotrope-like
10 compositions consist essentially of from about 84.5 to
about 98.9 weight percent 1,3-dichloro-1,1,2,2,3-
pentafluoropropane, from about 1 to about 6.5 weight
percent 2-propanol and from about 0.1 to about 9 weight
percent cyclohexane.

15 In another preferred embodiment of the invention
utilizing 225cb and cyclohexane, the azeotrope-like
compositions consist essentially of from about 83 to about
99.4 weight percent 1,3-dichloro-1,1,2,2,3-
20 pentafluoropropane, from about 0.5 to about 9 weight
percent 2-propanol and from about 0.1 to about 8 weight
percent cyclohexane.

In a more preferred embodiment of the invention
25 utilizing 225cb and cyclohexane, the azeotrope-like
compositions consist essentially of from about 85.5 to
about 98.9 weight percent 1,3-dichloro-1,1,2,2,3-
pentafluoropropane, from about 1 to about 6.5 weight
percent 2-propanol and from about 0.1 to about 8 weight
30 percent cyclohexane.

In a still more preferred embodiment of the
invention utilizing 225cb and cyclohexane, the
azeotrope-like compositions consist essentially of from
about 87 to about 98.4 weight percent
35 1,3-dichloro-1,1,2,2,3-pentafluoropropane, from about 1.5
to about 6 weight percent 2-propanol and from about 0.1 to
about 7 weight percent cyclohexane.

-13-

In a most preferred embodiment of the invention utilizing 225cb and cyclohexane, the azeotrope-like compositions consist essentially of from about 90.5 to about 97.5 weight percent 1,3-dichloro-1,1,2,2,3-pentafluoropropane, from about 1.5 to about 4.5 weight percent 2-propanol and from about 1 to about 5 weight percent cyclohexane.

10 The precise or true azeotrope compositions have not been determined but have been ascertained to be within the indicated ranges. Regardless of where the true azeotropes lie, all compositions within the indicated ranges, as well as certain compositions outside the indicated ranges, are
15 azeotrope-like, as defined more particularly below.

From fundamental principles, the thermodynamic state of a fluid is defined by four variables: pressure, temperature, liquid composition and vapor composition, or
20 P-T-X-Y, respectively. An azeotrope is a unique characteristic of a system of two or more components where X and Y are equal at a stated P and T. In practice, this means that the components of a mixture cannot be separated during distillation, and therefore are useful in vapor
25 phase solvent cleaning as described above.

For the purpose of this discussion, by azeotrope-like composition is intended to mean that the composition behaves like a true azeotrope in terms of its
30 constant-boiling characteristics or tendency not to fractionate upon boiling or evaporation. Such composition may or may not be a true azeotrope. Thus, in such compositions, the composition of the vapor formed during boiling or evaporation is identical or substantially
35 identical to the original liquid composition. Hence, during boiling or evaporation, the liquid composition, if it changes at all, changes only minimally. This is

-14-

contrasted with non-azeotrope-like compositions in which the liquid composition changes substantially during boiling a evaporation.

5 Thus, one way to determine whether a candidate mixture is "azeotrope-like" within the meaning of this invention, is to distill a sample thereof under conditions (i.e. resolution - number of plates) which would be
10 expected to separate the mixture into its separate components. If the mixture is non-azeotropic or non-azeotrope-like, the mixture will fractionate, i.e. separate into its various components with the lowest boiling component distilling off first, and so on. If the
15 mixture is azeotrope-like, some finite amount of a first distillation cut will be obtained which contains all of the mixture components and which is constant boiling or behaves as a single substance. This phenomenon cannot occur if the mixture is not azeotrope-like, i.e., it is
20 not part of an azeotropic system. If the degree of fractionation of the candidate mixture is unduly great, then a composition closer to the true azeotrope must be selected to minimize fractionation. Of course, upon
25 distillation of an azeotrope-like composition such as in a vapor degreaser, the true azeotrope will form and tend to concentrate.

 It follows from the above that another characteristic of azeotrope-like compositions is that
30 there is a range of compositions containing the same components in varying proportions which are azeotrope-like. All such compositions are intended to be covered by the term azeotrope-like as used herein. As an example, it is well known that at different pressures, the composition
35 of a given azeotrope will vary at least slightly as does the boiling point of the composition. Thus, an azeotrope of A and B represents a unique type of relationship having

-15-

a variable composition depending on temperature and/or pressure. Accordingly, another way of defining azeotrope-like within the meaning of the invention is to state that
5 such mixtures boil within about $\pm 3.9^{\circ}\text{C}$ (at 760 mm Hg) of the 52.2°C boiling point disclosed herein. As is readily understood by persons skilled in the art, the boiling point of the azeotrope will vary with the pressure.

10 In the process embodiment of the invention, the azeotrope-like compositions of the invention may be used to clean solid surfaces by treating said surfaces with said compositions in any manner well known in the art such as by dipping or spraying or use of conventional
15 degreasing apparatus.

As stated above, the azeotrope-like compositions discussed herein are useful as solvents for various cleaning applications including vapor degreasing,
20 defluxing, cold cleaning, dry cleaning, dewatering, decontamination, spot cleaning, aerosol propelled rework, extraction, particle removal, and surfactant cleaning applications. These azeotrope-like compositions are also useful as blowing agents, Rankine cycle and absorption
25 refrigerants, and power fluids.

The dichloropentafluoropropane, 2-propanol and C_6 hydrocarbon components of the invention are known materials. Preferably, they should be used in
30 sufficiently high purity so as to avoid the introduction of adverse influences upon the solvents or constant boiling properties of the system.

Commercially available 2-propanol and C_6 hydrocarbons may be used in the present invention. Most
35 dichloropentafluoropropane isomers, like the preferred HCFC-225ca isomer, however, are not available in

-16-

commercial quantities, therefore, until such time as they become commercially available, they may be prepared by following the organic syntheses disclosed herein. For example, 1,1-dichloro-2,2,3,3,3-pentafluoropropane, may be prepared by reacting 2,2,3,3,3-pentafluoro-1-propanol and p-toluenesulfonate chloride together to form 2,2,3,3,3-pentafluoropropyl-p-toluenesulfonate. Next, N-methylpyrrolidone, lithium chloride, and the 2,2,3,3,3-pentafluoropropyl-p-toluenesulfonate are reacted together to form 1-chloro-2,2,3,3,3-pentafluoropropane. Finally, chlorine and the 1-chloro-2,2,3,3,3-pentafluoropropane are reacted together to form 1,1-dichloro-2,2,3,3,3-pentafluoropropane. A detailed synthesis is set forth in Example 1.

Synthesis of 2,2-dichloro-1,1,1,3,3-pentafluoropropane (225a). This compound may be prepared by reacting a dimethylformamide solution of 1,1,1-trichloro-2,2,2-trifluoromethane with chlorotrimethylsilane in the presence of zinc, forming 1-(trimethylsiloxy)-2,2-dichloro-3,3,3-trifluoro-N,N-dimethylpropylamine. The 1-(trimethylsiloxy)-2,2-dichloro-3,3,3-trifluoro-N,N-dimethyl propylamine is reacted with sulfuric acid to form 2,2-dichloro-3,3,3-trifluoropropionaldehyde. The 2,2-dichloro-3,3,3-trifluoropropionaldehyde is then reacted with sulfur tetrafluoride to produce 2,2-dichloro-1,1,1,3,3-pentafluoropropane.

Synthesis of 1,2-dichloro-1,2,3,3,3-pentafluoropropane (225ba). This isomer may be prepared by the synthesis disclosed by O. Paleta et al., Bull. Soc. Chim. Fr., (6) 920-4 (1986).

Synthesis of 1,2-dichloro-1,1,2,3,3-pentafluoropropane (225bb). The synthesis of this isomer is disclosed by M. Hauptschein and L.A. Bigelow, J. Am. Chem.

-17-

Soc., (73) 1428-30 (1951). The synthesis of this compound is also disclosed by A.H. Fainberg and W.T. Miller, Jr., J. Am. Chem. Soc., (79) 4170-4, (1957).

5

Synthesis of 1,3-dichloro-1,1,2,2,3-pentafluoropropane (225cb). The synthesis of this compound involves four steps.

10 Part A - Synthesis of 2,2,3,3-tetrafluoropropyl-p-toluenesulfonate. 406 gm (3.08 mol) 2,2,3,3-tetrafluoropropanol, 613 gm (3.22 mol) tosylchloride, and 1200 ml water were heated to 50°C with mechanical stirring. Sodium hydroxide (139.7 gm, 3.5 ml) in 560 ml water was added at a rate such that the temperature remained less than 65°C. After the addition was completed, the
15 mixture was stirred at 50°C until the pH of the aqueous phase was 6. The mixture was cooled and extracted with 1.5 liters methylene chloride. The organic layer was washed twice with 200 ml aqueous ammonia, 350 ml water, dried with magnesium sulfate, and distilled to give
20 697.2 gm (79%) viscous oil.

Part B - Synthesis of 1,1,2,2,3-pentafluoropropane. A 500 ml flask was equipped with a mechanical stirrer and a Vigreux distillation column, which in turn
25 was connected to a dry-ice trap, and maintained under a nitrogen atmosphere. The flask was charged with 400 ml N-methylpyrrolidone, 145 gm (0.507 mol) 2,2,3,3-tetrafluoropropyl-p-toluenesulfonate (produced in Part A above), and 87 gm (1.5 mol) spray-dried KF. The mixture
30 was then heated to 190-200°C for about 3.25 hours during which time 61 gm volatile product distilled into the cold trap (90% crude yield). Upon distillation, the fraction boiling at 25-28°C was collected.

Part C - Synthesis of 1,1,3-trichloro-1,2,2,3,3-pentafluoropropane. A 22 liter flask was evacuated and
35 charged with 20.7 gm (0.154 mol) 1,1,2,2,3-pentafluoropropane (produced in Part B above) and 0.6 mol chlorine.

-18-

It was irradiated 100 minutes with a 450 W Hanovia Hg lamp at a distance of about 3 inches (7.6 cm). The flask was then cooled in an ice bath, nitrogen being added as
5 necessary to maintain 1 atm (101 kPa). Liquid in the flask was removed via syringe. The flask was connected to a dry-ice trap and evacuated slowly (15-30 minutes). The contents of the dry-ice trap and the initial liquid phase totaled 31.2 g (85%), the GC purity being 99.7%. The
10 product from several runs was combined and distilled to provide a material having b.p. 73.5-74°C.

Part D - Synthesis of 1,3-dichloro-1,1,2,2,3-pentafluoropropane. 106.6 gm (0.45 mol) of 1,1,3-trichloro-1,2,2,3,3-pentafluoropropane (produced in Part C
15 above) and 300 gm (5 mol) isopropanol were stirred under an inert atmosphere and irradiated 4.5 hours with a 450 W Hanovia Hg lamp at a distance of 2-3 inches (5-7.6 cm). The acidic reaction mixture was then poured into 1.5 liters ice water. The organic layer was separated, washed
20 twice with 50 ml water, dried with calcium sulfate, and distilled to give 50.5 gm $\text{ClCF}_2\text{CF}_2\text{CHClF}$, bp 54.5-56°C (55%). ^1H NMR (CDCl_3): ddd centered at 6.43 ppm. $J_{\text{H-C-F}} = 47$ Hz, $J_{\text{H-C-C-Fa}} = 12$ Hz, $J_{\text{H-C-C-Fb}} = 2$ Hz.

25

Synthesis of 1,1-dichloro-1,2,2,3,3-pentafluoropropane (225cc). This compound may be prepared by reacting 2,2,3,3-tetrafluoro-1-propanol and
p-toluenesulfonate chloride to form 2,2,3,3-tetrafluoropropyl-p-toluenesulfonate. Next, the 2,2,3,3-
30 tetrafluoropropyl-p-toluenesulfonate is reacted with potassium fluoride in N-methylpyrrolidone to form 1,1,2,2,3-pentafluoropropane. Then, the 1,1,2,2,3-pentafluoropropane is reacted with chlorine to form
35 1,1-dichloro-1,2,2,3,3-pentafluoropropane.

-19-

Synthesis of 1,2-dichloro-1,1,3,3,3-pentafluoropropane (225d). This isomer is commercially available from P.C.R. Incorporated of Gainesville, Florida.

5 Alternately, this compound may be prepared by adding equimolar amounts of 1,1,1,3,3-pentafluoropropane and chlorine gas to a borosilicate flask that has been purged of air. The flask is then irradiated with a mercury lamp. Upon completion of the irradiation, the contents of

10 the flask are cooled. The resulting product will be 1,2-dichloro-1,1,3,3,3-pentafluoropropane.

Synthesis of 1,3-dichloro-1,1,2,3,3-pentafluoropropane (225ea). This compound may be prepared by

15 reacting trifluoroethylene with dichlorotri-fluoromethane to produce 1,3-dichloro-1,1,2,3,3,-pentafluoropropane and 1,1-dichloro-1,2,3,3,3-pentafluoropropane. The 1,3-dichloro-1,1,2,3,3-pentafluoropropane is separated from its isomers using fractional

20 distillation and/or preparative gas chromatography.

Synthesis of 1,1-dichloro-1,2,3,3,3-pentafluoropropane (225eb). This compound may be prepared by reacting trifluoroethylene with dichlorodifluoromethane to

25 produce 1,3-dichloro-1,1,2,3,3-pentafluoropropane and 1,1-dichloro-1,2,3,3,3-pentafluoropropane. The 1,1-dichloro-1,2,3,3,3-pentafluoropropane is separated from its isomer using fractional distillation and/or preparative gas chromatography. Alternatively, 225eb may

30 be prepared by a synthesis disclosed by O. Paleta et al., Bull. Soc. Chim. Fr., (6) 920-4 (1986). The 1,1-dichloro-1,2,3,3,3-pentafluoropropane can be separated from its two isomers using fractional distillation and/or preparative gas chromatography.

35 It should be understood that the present compositions may include additional components which form

-20-

new azeotrope-like compositions. Any such compositions are considered to be within the scope of the present invention as long as the compositions are constant-boiling or essentially constant-boiling and contain all of the essential components described herein.

Inhibitors may be added to the present azeotrope-like compositions to inhibit decomposition of the compositions; react with undesirable decomposition products of the compositions; and/or prevent corrosion of metal surfaces. Any or all of the following classes of inhibitors may be employed in the invention: epoxy compounds such as propylene oxide; nitroalkanes such as nitromethane; ethers such as 1-4-dioxane; unsaturated compounds such as 1,4-butyne diol; acetals or ketals such as dipropoxy methane; ketones such as methyl ethyl ketone; alcohols such as tertiary amyl alcohol; esters such as triphenyl phosphite; and amines such as triethyl amine. Other suitable inhibitors will readily occur to those skilled in the art.

Having described the invention in detail and by reference to preferred embodiments thereof, it will be apparent that modifications and variations are possible without departing from the scope of the invention defined in the appended claims.

The present invention is more fully illustrated by the following non-limiting Examples.

Example 1

This example is directed to the preparation of 1,1-dichloro-2,2,3,3,3-pentafluoropropane (HCFC-225ca).

-21-

Part A - Synthesis of 2,2,3,3,3-pentafluoropropyl-p-toluenesulfonate. To p-toluenesulfonate chloride(400.66g, 2.10mol) in water at 25°C was added
5 2,2,3,3,3-pentafluoro-1-propanol (300.8g). The mixture was heated in a 5 liter, 3-neck separatory funnel type reaction flask, under mechanical stirring, to a temperature of 50°C. Sodium hydroxide(92.56g, 2.31mol) in
10 383ml water(6M solution) was added dropwise to the reaction mixture via addition funnel over a period of 2.5 hours, keeping the temperature below 55°C. Upon completion of this addition, when the pH of the aqueous phase was approximately 6, the organic phase was drained from the flask while still warm, and allowed to cool to
15 25°C. The crude product was recrystallized from petroleum ether to afford white needles of 2,2,3,3,3-pentafluoropropyl-p-toluenesulfonate(500.7g, 1.65mol, 82.3%).

Part B - Synthesis of 1-chloro-2,2,3,3,3-pentafluoropropane. A 1 liter flask fitted with a
20 thermometer, Vigreux column and distillation receiving head was charged with 248.5g(0.82mol) 2,2,3,3,3-pentafluoropropyl-p-toluenesulfonate(produced in Part A above), 375ml N-methylpyrrolidone, and 46.7 g(1.1mol)
25 lithium chloride. The mixture was then heated with stirring to 140°C at which point, product began to distill over. Stirring and heating were continued until a pot temperature of 198°C had been reached at which point, there was no further distillate being collected. The
30 crude product was re-distilled to give 107.2g(78%) of product.

Part C - Synthesis of 1,1-dichloro-2,2,3,3,3-pentafluoropropane. Chlorine (289ml/min) and
35 1-chloro-2,2,3,3,3-pentafluoropropane(produced in Part B above) (1.72g/min) were fed simultaneously into a 1 inch (2.54cm) x 2 inches(5.08cm) monel reactor at 300°C. The process was repeated until 184g crude product had

-22-

collected in the cold traps exiting the reactor. After the crude product was washed with 6 M sodium hydroxide and dried with sodium sulfate, it was distilled to give 69.2g starting material and 46.8g 1,1,-dichloro-2,2,3,3,3-pentafluoropropane (bp 48-50.5°C). ¹H NMR: 5.9 (t, J=7.5 H) ppm; ¹⁹F NMR: 79.4 (3F) and 119.8 (2F) ppm upfield from CFC1₃.

10

Examples 2-8

The compositional range over which 225ca, 2-propanol and n-hexane exhibit constant-boiling behavior was determined. This was accomplished by charging selected 225ca-based binary compositions into an ebulliometer, bringing them to a boil, adding measured amounts of a third component and finally recording the temperature of the ensuing boiling mixture. In each case, a minimum in the boiling point versus composition curve occurred; indicating that a constant boiling composition formed.

The ebulliometer consisted of a heated sump in which the 225ca-based binary mixture was brought to a boil. The upper part of the ebulliometer connected to the sump was cooled thereby acting as a condenser for the boiling vapors, allowing the system to operate at total reflux. After bringing the 225ca-based binary mixture to boil at atmospheric pressure, measured amounts of a third component were titrated into the ebulliometer. The change in boiling point was measured with a platinum resistance thermometer.

To normalize observed boiling points during different days to 760 millimeters of mercury pressure, the approximate normal boiling points of 225ca-based mixtures were estimated by applying a barometric correction factor of about 26 mm Hg/°C, to the observed values. However, it

-23-

is to be noted that this corrected boiling point is generally accurate up to $\pm 0.4^{\circ}\text{C}$ and serves only as a rough comparison of boiling points determined on different days.

The following table lists, for Examples 2-8, the compositional range over which the 225ca/2-propanol/n-hexane mixture is constant boiling, i.e., the boiling point deviations are within $\pm 0.5^{\circ}\text{C}$ of each other. Based on the data in Table I, compositions of 225ca/2-propanol/n-hexane ranging from about 87-99.8/0.1-4/0.1-9 and preferably 89.5-99.4/0.1-2.5/0.5-8 weight percent respectively would exhibit constant boiling behavior.

TABLE I

<u>Example</u>		<u>Starting Binary Composition (Wt%)</u>		
20	2	225ca/2-propanol (99/1)		
	3	225ca/2-propanol (99.5/0.5)		
	4	225ca/n-hexane(94.5/5.5)		
	5	225ca/n-hexane(96.5/3.5)		
	6	225ca/n-hexane (98/2)		
25 <u>Example</u>		<u>Range over which third component is constant boiling</u>	<u>Preferred Range</u>	<u>Minimum Temp. ($^{\circ}\text{C}$)</u>
30	2	0.1-9.5 n-hex.	0.5-8.0 n-hex.	51.2
	3	0.1-9.0 n-hex.	0.5-7.5 n-hex.	51.2
	4	0.1-3.2 2-prop.	0.1-2.1 2-prop.	51.1
	5	0.1-4.4 2-prop.	0.1-2.5 2-prop.	51.2
	6	0.1-2.8 2-prop.	0.1-2.0 2-prop.	51.1

Examples 7-13

The compositional range over which 225cb, 2-propanol and cyclohexane exhibit constant-boiling behavior was determined. This was accomplished by repeating the experiment outlined in Examples 2-6 above except that 225cb is substituted for 225ca and cyclohexane was used in place of n-hexane.

-24-

Table II lists the compositional range over which the 225cb/2-propanol/cyclohexane mixture is constant boiling, i.e., the boiling point deviations are within $\pm 0.5^{\circ}\text{C}$ of each other. Based on the data in Table II, compositions of 225cb/2-propanol/cyclohexane ranging from about 82-99.4/0.5-9/0.1-9 weight percent and preferably about 87-98.4/1.5-6/0.1-7 weight percent respectively would exhibit constant boiling behavior.

10

TABLE II

<u>Example</u>		<u>Starting Binary Composition (wt %)</u>		
15	7	225cb/2-propanol (96.5/3.5)		
	8	225cb/2-propanol (98/2)		
	9	225cb/2-propanol (95.5/4.5)		
	10	225cb/cyclohexane (97.2/2.8)		
	11	225cb/cyclohexane (97.9/2.1)		
	12	225cb/cyclohexane (94.5/5.5)		
	13	225cb/cyclohexane (99/1)		
<u>Example</u>		<u>Range over which third component is constant boiling</u>	<u>Preferred Range</u>	<u>Minimum Temp. ($^{\circ}\text{C}$)</u>
20	7	0.1-9.0 cyclohex.	0.1-8 cyclohex.	55.6
	8	0.1-7.0 cyclohex.	0.1-7 cyclohex.	55.6
	9	0.1-8.5 cyclohex.	0.1-6.5 cyclohex.	55.6
	10	0.5-6.5 2-prop.	1.0-5 2-prop.	55.7
	11	0.5-9.2 2-prop.	1.3-6.5 2-prop.	55.6
	12	0.5-10.0 2-prop.	1.3-6.5 2-prop.	55.7
	13	0.5-8.5 2-prop.	1.0-5.8 2-prop.	55.6

Examples 14-25

The azeotropic properties of the below listed dichloropentafluoropropane components (Table III) with 2-propanol and n-hexane are studied. This is accomplished by charging selected dichloropentafluoropropane-based binary compositions into an ebulliometer, bringing them to a boil, adding measured amounts of a third component and finally recording the temperature of the ensuing boiling mixture. In each case a minimum in the boiling point versus composition curve occurs, indicating that a

-25-

constant boiling composition forms between each dichloropentafluoropropane component, 2-propanol and n-hexane.

5

TABLE IIIDichloropentafluoropropane Component

10	2,2-dichloro-1,1,1,3,3-pentafluoropropane (225a)
	1,2-dichloro-1,2,3,3,3-pentafluoropropane (225ba)
	1,2-dichloro-1,1,2,3,3-pentafluoropropane (225bb)
	1,3-dichloro-1,1,2,2,3-pentafluoropropane (225cb)
15	1,1-dichloro-1,2,2,3,3-pentafluoropropane (225cc)
	1,2-dichloro-1,1,3,3,3-pentafluoropropane (225d)
	1,3-dichloro-1,1,2,3,3-pentafluoropropane (225ea)
	1,1-dichloro-1,2,3,3,3-pentafluoropropane (225eb)
20	1,1-dichloro-2,2,3,3,3-pentafluoropropane/(mixture of 1,3-dichloro-1,1,2,2,3-pentafluoropropane 225ca/cb)
	1,1-dichloro-1,2,2,3,3-pentafluoropropane/(mixture of 1,3-dichloro-1,1,2,2,3-pentafluoropropane 225eb/cb)

Examples 26-35

25

The azeotropic properties of the below-listed dichloropentafluoropropane components (Table IV) with 2-propanol and cyclohexane are studied by repeating the experiment outlined in Examples 14-25 above except that

30 cyclohexane is substituted for n-hexane. In each case, a minimum in the boiling point versus composition curve occurs indicating that a constant boiling composition forms between each dichloropentafluoropropane component, 2-propanol and cyclohexane.

35

-26-

TABLE IV

	<u>Dichloropentafluoropropane Component</u>
5	2,2-dichloro-1,1,1,3,3-pentafluoropropane (225a)
	1,2-dichloro-1,2,3,3,3-pentafluoropropane (225ba)
	1,2-dichloro-1,1,2,3,3-pentafluoropropane (225bb)
10	1,1-dichloro-2,2,3,3,3-pentafluoropropane (225ca)
	1,1-dichloro-1,2,2,3,3-pentafluoropropane (225cc)
	1,2-dichloro-1,1,3,3,3-pentafluoropropane (225d)
	1,3-dichloro-1,1,2,3,3-pentafluoropropane (225ea)
15	1,1-dichloro-1,2,3,3,3-pentafluoropropane (225eb)
	1,1-dichloro-2,2,3,3,3-pentafluoropropane/(mixture of 1,3-dichloro-1,1,2,2,3-pentafluoropropane 225ca/cb)
	1,1-dichloro-1,2,2,3,3-pentafluoropropane/(mixture of 1,3-dichloro-1,1,2,2,3-pentafluoropropane 225eb/cb)
20	<u>Examples 36-46</u>

The azeotropic properties of the below-listed dichloropentafluoropropane components (Table V) with 2-propanol and 2-methylpentane are studied by repeating the experiment outlined in Examples 14-25 above except that 2-methylpentane is substituted for n-hexane. In each case, a minimum in the boiling point versus composition curve occurs indicating that a constant boiling composition forms between each dichloropentafluoropropane component, 2-propanol and 2-methylpentane.

TABLE V

	<u>Dichloropentafluoropropane Component</u>
35	2,2-dichloro-1,1,1,3,3-pentafluoropropane (225a)
	1,2-dichloro-1,2,3,3,3-pentafluoropropane (225ba)

-27-

- 1,2-dichloro-1,1,2,3,3-pentafluoropropane (225bb)
1,1-dichloro-2,2,3,3,3-pentafluoropropane (225ca)
1,3-dichloro-2,2,3,3,3-pentafluoropropane (225cb)
5 1,1-dichloro-1,2,2,3,3-pentafluoropropane (225cc)
1,2-dichloro-1,1,3,3,3-pentafluoropropane (225d)
1,3-dichloro-1,1,2,3,3-pentafluoropropane (225ea)
10 1,1-dichloro-1,2,3,3,3-pentafluoropropane (225eb)
1,1-dichloro-2,2,3,3,3-pentafluoropropane/(mixture of
1,3-dichloro-1,1,2,2,3-pentafluoropropane 225ca/cb)
1,1-dichloro-1,2,2,3,3-pentafluoropropane/(mixture of
1,3-dichloro-1,1,2,2,3-pentafluoropropane 225eb/cb)

15

Examples 47-57

- The azeotropic properties of the dichloropenta-
fluoropropane components listed in Table V with 2-propanol
and 3-methylpentane are studied by repeating the
20 experiment outlined in Examples 14-25 above except that
3-methylpentane is substituted for n-hexane. In each
case, a minimum in the boiling point versus composition
curve occurs indicating that a constant boiling
composition forms between each dichloropenta-
25 fluoropropane component, 2-propanol and 3-methylpentane.

Examples 58-68

- The azeotropic properties of the dichloropenta-
30 fluoropropane components listed in Table V with 2-propanol
and 2,2-dimethylbutane are studied by repeating the
experiment outlined in Examples 14-25 above except that
2,2-dimethylbutane is substituted for n-hexane. In each
case, a minimum in the boiling point versus composition
35 curve occurs indicating that a constant boiling
composition forms between each dichloropentafluoropropane
component, 2-propanol and 2,2-dimethylbutane.

-28-

Examples 69-79

5 The azeotropic properties of the dichloropenta-
fluoropropane components listed in Table V with 2-propanol
and 2,3-dimethylbutane are studied by repeating the
experiment outlined in Examples 14-25 above except that
2,3-dimethylbutane is substituted for n-hexane. In each
case, a minimum in the boiling point versus composition
10 curve occurs indicating that a constant boiling
composition forms between each dichloropentafluoropropane
component, 2-propanol and 2,3-dimethylbutane.

Examples 80-90

15 The azeotropic properties of the dichloropenta-
fluoropropane components listed in Table V with 2-propanol
and methylcyclopentane are studied by repeating the
experiment outlined in Examples 14-25 above except that
methylcyclopentane is substituted for n-hexane. In each
20 case, a minimum in the boiling point versus composition
curve occurs indicating that a constant boiling
composition forms between each dichloropentafluoropropane
component, 2-propanol and methylcyclopentane.

25

Examples 91-101

The azeotropic properties of the dichloropenta-
fluoropropane components listed in Table V with 2-propanol
and commercial isohexane grade 1 are studied by repeating
30 the experiment outlined in Examples 14-25 above except
that commercial isohexane grade 1 is substituted for
n-hexane. In each case, a minimum in the boiling point
versus composition curve occurs indicating that a constant
boiling composition forms between each dichloropentafluoro-
35 propane component, 2-propanol and commercial isohexane
grade 1.

-29-

Examples 102-112

The azeotropic properties of the dichloropenta-
5 fluoropropane components listed in Table V with 2-propanol
and commercial isohexane grade 2 are studied by repeating
the experiment outlined in Examples 14-25 above except
that commercial isohexane grade 2 is substituted for
n-hexane. In each case, a minimum in the boiling point
10 versus composition curve occurs indicating that a constant
boiling composition forms between each dichloropentafluoro-
propane component, 2-propanol and commercial isohexane
grade 2.

15

20

25

30

35

-30-

What is claimed is:

1. Azeotrope-like compositions comprising from
5 about 62 to about 99.8 weight percent dichloro-
pentafluoropropane, from about 0.1 to about 10 weight
percent 2-propanol and from about 0.1 to about 28 weight
percent C₆ hydrocarbon which boil at about 52.2°C at
760 mm Hg.
- 10 2. The azeotrope-like compositions of claim 1
wherein said compositions boil at about 52.2°C ± about
3.9°C at 760 mm Hg.
- 15 3. The azeotrope-like compositions of claim 1
wherein said compositions boil at about 52.4°C ± about
3.7°C at 760 mm Hg.
- 20 4. The azeotrope-like compositions of claim 1
wherein said compositions consist essentially of from
about 67 to about 96.5 weight percent dichloropenta-
fluoropropane, from about 0.5 to about 7 weight percent
2-propanol, and from about 3 to about 26 weight percent
C₆ hydrocarbon.
- 25 5. The azeotrope-like compositions of claim 1
wherein said compositions consist essentially of from
about 73 to about 95.5 weight percent dichloropenta-
fluoropropane, from about 0.5 to about 7 weight percent
2-propanol and from about 4 to about 20 weight percent
30 C₆ hydrocarbon.
- 35 6. The azeotrope-like compositions of claim 1
wherein said compositions consist essentially of from
about 83 to about 95.5 weight percent dichloropenta-
fluoropropane, from about 0.5 to about 7 weight percent
2-propanol and from about 4 to about 10 weight percent
C₆ hydrocarbon.

-31-

7. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 67 to about 89.5 weight percent dichloropenta-
5 fluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 10 to about 26 weight percent C_6 hydrocarbon

8. The azeotrope-like compositions of claim 1
10 wherein said compositions consist essentially of from about 73 to about 96.5 weight percent dichloropenta-fluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 3 to about 20 weight percent n-hexane and boil at about $52.6^{\circ}C$ at 760 mm Hg.

15 9. The azeotrope-like compositions of claim 8 wherein said compositions boil at about $52.6^{\circ}C \pm$ about $3.0^{\circ}C$ at 760 mm Hg.

20 10. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 67 to about 93.5 weight percent dichloropenta-fluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 6 to about 26 weight percent
25 2-methylpentane and boil at about $51.4^{\circ}C \pm$ about $2.5^{\circ}C$ at 760 mm Hg.

11. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from
30 about 71 to about 96.5 weight percent dichloropenta-fluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 3 to about 22 weight percent 3-methylpentane and boil at about $51.9^{\circ}C$ at 760 mm Hg.

35 12. The azeotrope-like compositions of claim 11 wherein said compositions boil at about $51.9^{\circ}C \pm$ about $2.9^{\circ}C$ at 760 mm Hg.

-32-

13. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 71 to about 96.5 weight percent dichloropenta-
5 fluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 3 to about 22 weight percent methylcyclopentane and boil at about 52.8°C at 760 mm Hg.

14. The azeotrope-like compositions of claim 13
10 wherein said compositions boil at about 52.8°C ± about 2.8°C at 760 mm Hg.

15. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from
15 about 67 to about 93.5 weight percent dichloropenta-fluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 6 to about 26 weight percent commercial isohexane grade 1 and boil at about 51.1°C at 760 mm Hg.

20 16. The azeotrope-like compositions of claim 15 wherein said compositions boil at about 51.1°C ± about 2.7°C at 760 mm Hg.

25 17. The azeotrope-like compositions of claim 15 wherein said compositions boil at about 51.1°C ± about 2.3°C at 760 mm Hg.

18. The azeotrope-like compositions of claim 1
30 wherein said compositions consist essentially of from about 67 to about 93.5 weight percent dichloropenta-fluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 6 to about 26 weight percent commercial isohexane grade 2 and boil at about 51.1°C at
35 760 mm Hg.

-33-

19. The azeotrope-like compositions of claim 17 wherein said compositions boil at about $51.1^{\circ}\text{C} \pm$ about 2.7°C at 760 mm Hg.

5

20. The azeotrope-like compositions of claim 17 wherein said compositions boil at about $51.1^{\circ}\text{C} \pm$ about 2.3°C at 760 mm Hg.

10

21. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 83 to about 99.4 weight percent dichloropentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 0.1 to about 10 weight percent cyclohexane and boil at about 53.3°C at 760 mm Hg.

15

22. The azeotrope-like compositions of claim 21 wherein said compositions boil at about $53.3^{\circ}\text{C} \pm$ about 2.8°C at 760 mm Hg.

20

23. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 87 to about 99.8 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 4 weight percent 2-propanol and from about 0.1 to about 9 weight percent n-hexane and boil at about 51.2°C at 760 mm Hg.

25

24. The azeotrope-like compositions of claim 23 wherein said compositions boil at about $51.2^{\circ}\text{C} \pm$ about 0.7°C at 760 mm Hg.

30

25. The azeotrope-like compositions of claim 23 wherein said compositions boil at about $51.2^{\circ}\text{C} \pm$ about 0.5°C at 760 mm Hg.

35

-34-

26. The azeotrope-like compositions of claim 23 wherein said compositions consist essentially of from about 88.5 to about 99.8 weight percent 1,1-dichloro-
5 2,2,3,3,3-pentafluoropropane, from about 0.1 to about 2.5 weight percent 2-propanol and from about 0.1 to about 9 weight percent n-hexane.

27. The azeotrope-like compositions of claim 23
10 wherein said compositions consist essentially of from about 88 to about 99.4 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 4 weight percent 2-propanol and from about 0.5 to about 8 weight percent n-hexane.

15 28. The azeotrope-like compositions of claim 23 wherein said compositions consist essentially of from about 89.5 to about 99.4 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 2.5
20 weight percent 2-propanol and from about 0.5 to about 8 weight percent n-hexane.

29. The azeotrope-like compositions of claim 23 wherein said compositions consist essentially of from
25 about 91 to about 99.4 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 2 weight percent 2-propanol and from about 0.5 to about 7 weight percent n-hexane.

30 30. The azeotrope-like compositions of claim 23 wherein said compositions consist essentially of from about 92 to about 98.9 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.1 to about 2
weight percent 2-propanol and from about 1 to about 6
35 weight percent n-hexane.

-35-

31. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 70 to about 92.5 weight percent 1,1-dichloro-
5 2,2,3,3,3-pentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 7 to about 23 weight percent 2-methylpentane and boil at about 49.7°C at 760 mm Hg.

10 32. The azeotrope-like compositions of claim 31 wherein said compositions boil at about 49.7°C \pm about 0.9°C at 760 mm Hg.

15 33. The azeotrope-like compositions of claim 31 wherein said compositions boil at about 49.7°C \pm about 0.5°C at 760 mm Hg.

20 34. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 68 to about 92.5 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 7 to about 25 weight percent commercial isohexane grade 1 and boil at about 49.6°C at 760 mm Hg.

25 35. The azeotrope-like compositions of claim 34 wherein said compositions boil at about 49.6°C \pm about 1.3°C at 760 mm Hg.

30 36. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 68 to about 92.5 weight percent 1,1-dichloro-2,2,3,3,3-pentafluoropropane, from about 0.5 to about 7 weight percent 2-propanol and from about 7 to about 25 weight percent commercial isohexane grade 2 and boil at
35 about 49.6°C at 760 mm Hg.

-36-

37. The azeotrope-like compositions of claim 36 wherein said compositions boil at about 49.6°C \pm about 1.3°C at 760 mm Hg.

5

38. The azeotrope-like compositions of claim 1 wherein said compositions consist essentially of from about 82 to about 99.4 weight percent 1,3-dichloro-1,1,2,2,3-pentafluoropropane, from about 0.5 to about 9 weight percent 2-propanol and from about 0.1 to about 9 weight percent cyclohexane and boil at about 55.6°C at 760 mm Hg.

10

39. The azeotrope-like compositions of claim 38 wherein said compositions boil at about 55.6°C \pm about 0.5°C at 760 mm Hg.

15

40. The azeotrope-like compositions of claim 38 wherein said compositions boil at about 55.6°C \pm about 0.2°C at 760 mm Hg.

20

41. The azeotrope-like compositions of claim 38 wherein said compositions consist essentially of from about 84.5 to about 98.9 weight percent 1,3-dichloro-1,1,2,2,3-pentafluoropropane, from about 1 to about 6.5 weight percent 2-propanol and from about 0.1 to about 9 weight percent cyclohexane.

25

42. The azeotrope-like compositions of claim 38 wherein said compositions consist essentially of from about 83 to about 99.4 weight percent 1,3-dichloro-1,1,2,2,3-pentafluoropropane, from about 0.5 to about 9 weight percent 2-propanol and from about 0.1 to about 8 weight percent cyclohexane.

30

43. The azeotrope-like compositions of claim 38 wherein said compositions consist essentially of from about 85.5 to about 98.9 weight percent 1,3-dichloro-

35

-37-

1,1,2,2,3-pentafluoropropane, from about 1 to about 6.5 weight percent 2-propanol and from about 0.1 to about 8 weight percent cyclohexane.

5

44. The azeotrope-like compositions of claim 38 wherein said compositions consist essentially of from about 87 to about 98.4 weight percent 1,3-dichloro-1,1,2,2,3-pentafluoropropane, from about 1.5 to about 6 weight percent 2-propanol and from about 0.1 to about 7 weight percent cyclohexane.

10

45. The azeotrope-like compositions of claim 38 wherein said compositions consist essentially of from about 90.5 to about 97.5 weight percent 1,3-dichloro-1,1,2,2,3-pentafluoropropane, from about 1.5 to about 4.5 weight percent 2-propanol and from about 1 to about 5 weight percent cyclohexane.

15

46. The azeotrope-like compositions of claim 1 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

20

47. The azeotrope-like compositions of claim 5 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

25

48. The azeotrope-like compositions of claim 7 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

30

49. The azeotrope-like compositions of claim 8 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

35

-38-

50. The azeotrope-like compositions of claim 10 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and
5 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

51. The azeotrope-like compositions of claim 11 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and
10 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

52. The azeotrope-like compositions of claim 13 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and
15 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

53. The azeotrope-like compositions of claim 15 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and
20 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

54. The azeotrope-like compositions of claim 18 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and
25 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

55. The azeotrope-like compositions of claim 21 wherein said dichloropentafluoropropane is a mixture of 1,1-dichloro-2,2,3,3,3-pentafluoropropane and
30 1,3-dichloro-1,1,2,2,3-pentafluoropropane.

56. The azeotrope-like compositions of claim 1 wherein an effective amount of an inhibitor is optionally present in said composition.

35 57. The azeotrope-like compositions of claim 29 wherein an effective amount of an inhibitor is optionally present in said composition.